



Human Biology Review (ISSN 22774424)

www.HumanBiologyJournal.com

International Peer Reviewed Journal of Biological Anthropology

Volume 12, Number 4, October –December 2023 Issue

Original scientific paper

Projection of Life Expectancy at Birth for the Population of Urban Area in Bangladesh

M.S. Hossain¹, M.R. Islam^{2*}, M. K. Ali³, M.M. Rana⁴ and M. Shahiduzzaman⁵

Human Biology Review, 12 (4), pp. 235-251

Revised and accepted on August 30, 2023

1Md. Sabbir Hossain, Assistant Professor (Mathematics), Department of Business Administration, Bangladesh Islami University, Dhaka, Bangladesh, E-mail: sabbir169@yahoo.com

2Md. Rafiqul Islam, Professor, Department of Population Science and Human Resource Development, University of Rajshahi, Bangladesh, E-mail: rafique_pops@yahoo.com

3M. Korban Ali

Retired Professor, Department of Population Science and Human Resource Development, University of Rajshahi, Bangladesh, E-mail: Korban48@yahoo.com

4Md. Masud Rana, Institute of Water and Flood Management (IWF), Bangladesh University of Engineering and Technology (BUET), Dhaka 1000, Bangladesh. E-mail: ma-sud.raj82@gmail.com

5Md. Shahiduzzaman

Associate Professor, Department of Electrical and Electronic Engineering, Northern University Bangladesh, Dhaka-1230, Bangladesh. E-mail: shahidnub@yahoo.com

*Corresponding Author: Md. Rafiqul Islam, Professor,

Department of Population Science and Human Resource Development, University of Rajshahi, Bangladesh, E-mail: rafique_pops@yahoo.com

Projection of Life Expectancy at Birth for the Population of Urban Area in Bangladesh

M.S. Hossain, M.R. Islam, M. K. Ali, M.M. Rana and M. Shahiduzzaman

Citation: Hossain MS, Islam MR, Ali MK, Rana MM and Shahiduzzaman M. 2023. Projection of Life Expectancy at Birth for the Population of Urban Area in Bangladesh. Human Biology Review, 12 (4), 235-251.

ABSTRACT

Background: Life expectancy at birth (LE) reflects the overall mortality of a population and it is also used as one of the indicators of the health status of population.

Objective: The main purpose of this study is to project LE for the population of male, female and both sexes of urban region in Bangladesh by using polynomial model.

Data and Methodology: For this purpose, the data is obtained from Statistical Year Book (SYB) of Bangladesh of 2012. Quasi Newton Method is used to build up these models using the Statistical software STATISTICA. Furthermore, *t*-test statistics, *F*-test statistics and cross validity prediction power (CVPP) are applied to identify the correctness of these model.

Results: It is found that LE for the population of urban region of Bangladesh follow simple linear regression model. Based on *t*-stats, coefficient of determination, shrinkage coefficients and *F*-stats these models are seemed to be well fitted. So, these models can be used for better prediction. Then, these LE are projected during 2012-2051 using the fitted time trend models. It is observed that the LE for the population of male, female and both sexes of urban region of Bangladesh in 2051 are 75, 84.1 and 78 years respectively.

Conclusion: Urban population of Bangladesh are going to be aged population dramatically. Proper planning for the allocation of public and private resources may be taken for the better lifestyle of urban population.

Keywords: Life expectancy at birth (LE), Polynomial model, *t*-test statistics, *F*-test statistics, Cross validity prediction power (CVPP), Bangladesh

INTRODUCTION

Life expectancy is demarcated as the average number of supplementary years that a person of a specific age group will live, if the current age-specific mortality schedules remain constant over the course of the individual's lifetime (Chiang, 1968). Life expectancy is an indirect measure of mortality and it is also the main indicator of health status of a society or a country or a region. On the other hand, education and health are the leading sectors for socio-economic development for the developing countries like Bangladesh. Life expectancies in urban and rural areas for the population of male, female and both sexes from 1994 to 2011 show the shortest life expectancy of those living in rural area than that of urban areas (BBS, 2014). Life expectancy at birth (LE) in Bangladesh is poor in international comparisons with median of LE for male and female are 70 and 76 years respectively in 2011 (WHO, 2013). Life expectancy is not the measure of the aged population, but it shows the present situation of socio-economic development of a country. Moreover, it is found that LE of Bangladesh is increasing with the increase of time (BBS, 2013). Like many other nations, in Bangladesh female live lengthier than male and this gender difference in the LE has been increasing with passing of time. The longer LE of female population does not necessarily provide that the female are healthier than male. Studies on self-reported health status measure specify higher prevalence of functional restrictions and poor health among female (Parahyba et al., 2005; Lima et al., 2009; Zunzunegui et al., 2009 and Szwarcwald et al., 2011), suggesting that the additional years cannot essentially be lived in hale and hearty life. LE is increasing every year that means aged population is increasing in Bangladesh. So, especially government and policy makers should have clear concept about the number of aged population to ensure them with healthy environment. For this, correct information of LE in the future years is needed to take several steps for development of education, health, environment and others socioeconomic related sectors of Bangladesh. For this reason, mathematical model is selected to observe the trend of LE for the population of urban areas in Bangladesh. To do so, polynomial model has been chosen in this study. It is mentioned here that polynomial model was applied in a number of studies like as (Islam, 2004, 2005a; 2005b; 2007a and 2011; Islam et al., 2004 and 2014; Islam and Hossain, 2014a and 2014b). Therefore, the fundamental aims of this study are to construct some models to LE for the population of male, female and both sexes of urban region in

Bangladesh and then to project LE for them employing these fitted models for the period 2012-2051.

SOURCE OF DATA

A data on LE for the population of men, women and both sexes of urban region in Bangladesh during 2002-2011 have been obtained from SYB of Bangladesh of 2012 (BBS, 2013) which are shown in Table 1. To fulfill the above objectives, these LE data are used in this study.

Table 1: Observed LE for Male, Female and Both Sexes of the Population of Urban Area in Bangladesh during 2002-2011

| Year | Male | Female | Both Sexes |
|------|------|--------|------------|
| 2002 | 67.0 | 67.3 | 67.2 |
| 2003 | 67.3 | 67.9 | 67.6 |
| 2004 | 67.5 | 68.1 | 67.8 |
| 2005 | 67.6 | 68.1 | 67.9 |
| 2006 | 67.7 | 68.7 | 68.1 |
| 2007 | 67.7 | 68.7 | 68.1 |
| 2008 | 67.9 | 68.8 | 68.3 |
| 2009 | 68.2 | 69.2 | 68.7 |
| 2010 | 68.3 | 69.5 | 68.9 |
| 2011 | 68.9 | 71.1 | 69.9 |

Source: BBS (2013) Statistical Year Book of Bangladesh 2012.

METHODS

Data Smoothing

In this paper, it is found that there are some kinds of unexpected distortions in the data set if LE is dotted on a graph paper. For that reason, an adjustment is imperative and needed to abate these unexpected distortions before going to construct model to these data. In this situation, the data of LE for the population of men, women and both sexes of are smoothed by the Package Minitab Release 12.1 by smoothing technique named “4253H, twice” (Velleman, 1980). After that, these smoothed data are taken to build models and these smoothed data are presented in the Table 2.

Table 2: Smoothed and Predicted LE for Male, Female and Both Sexes of the Population of Urban Area in Bangladesh during 2002-2011

| Year | Male | | Female | | Both Sexes | |
|------|----------|-----------|----------|-----------|------------|-----------|
| | Smoothed | Predicted | Smoothed | Predicted | Smoothed | Predicted |
| 2002 | 67.0 | 67.0 | 67.4 | 67.3 | 67.2 | 67.2 |
| 2003 | 67.3 | 67.2 | 67.8 | 67.6 | 67.6 | 67.4 |
| 2004 | 67.5 | 67.3 | 68.1 | 68.0 | 67.8 | 67.7 |
| 2005 | 67.6 | 67.5 | 68.3 | 68.3 | 67.9 | 67.9 |
| 2006 | 67.7 | 67.6 | 68.5 | 68.7 | 68.0 | 68.1 |
| 2007 | 67.7 | 67.8 | 68.7 | 69.0 | 68.1 | 68.3 |
| 2008 | 67.9 | 68.0 | 69.1 | 69.3 | 68.3 | 68.5 |
| 2009 | 68.1 | 68.1 | 69.5 | 69.7 | 68.6 | 68.8 |
| 2010 | 68.4 | 68.3 | 69.9 | 70.0 | 69.0 | 69.0 |
| 2011 | 68.7 | 68.5 | 71.0 | 70.4 | 69.6 | 69.2 |

Model Building

The polynomial model is chosen to fit model (using the scattered plot of Fig.1-Fig.3) to LE for the population of men, women and both sexes in urban region of Bangladesh. Therefore, the configuration of n degree polynomial model is addressed by

$$y_t = a_0 + \sum_{i=1}^n a_i t^i + u$$

where, t represents time (years); y_t represents LE; the coefficient of t^i is a_i , a_0 is the constant term and u is the error term of the model. In this case, n is chosen so that the error sum of square is minimum. Moreover, it is noted that actually disturbance term (u) of the model is normally as well as independently distributed with mean zero and constant variance, that is, $u_i \sim \text{NID}(0, \sigma^2)$ (Gujarati, 1998). In this case, it is found that if $n=1$, i. e., the polynomial model of degree one becomes simple linear regression model fitted to these data sets. It is noted that these models are built using the Statistical software STATISTICA.

Model Validation Procedure

To verify how a large amount of these models on LE are stable, the cross validity prediction power (CVPP), ρ_{cv}^2 , is used for the urban area of Bangladesh. Here

$$\rho_{cv}^2 = 1 - \frac{(n-1)(n-2)(n+1)}{n(n-k-1)(n-k-2)}(1-R^2); \text{ where, } n \text{ indicates the number of cases, } k$$

specifies the number of predictors for the model and the cross validated R denotes the correlation between observed and predicted values of the response variable (Stevens, 1996). The positive value of $(\rho_{cv}^2 - R^2)$ mentions the shrinkage coefficient of the model; where ρ_{cv}^2 is CVPP and R^2 is the coefficient of determination of the model as well as the stability of R^2 of the model is (1-shrinkage). The results on model fittings and estimated CVPP are presented in Table 3. A number of studies have been used this technique as validation (Islam, 2006a; 2006b; 2006c; 2007b; 2007c; 2008; 2012a; 2012b; 2013; Islam & Hossain, 2013a; 2013b; Hossain & Islam, 2013; Islam et al., 2013; Islam and Hoque, 2015; Islam et al., 2020; Islam and Shitan, 2022).

F-test statistics

The F-test is employed to corroborate the measurement of overall significance of the model. The mathematical shape for F-test is specified under:

$$F = \frac{R^2 / (k-1)}{(1-R^2) / (n-k)}$$

where k represents the number of parameters of the model, n is the number of cases and R^2 is the coefficient of determination of the fitted model (Gujarati, 1998). These results are exposed in Table 4.

RESULTS

Table 1 and Table 2 represent the observed, smoothed and predicted LE for the population of men, women and both sexes of urban area in Bangladesh during 2002-2011. To see the level and trend of observed LE for the population, the data have been presented in graph paper shown in Fig. 4. It is investigated that the trend of LE for the population of men, women and both sexes are showing upward. It is also seen that LE for female is higher than male during 2002-2011. Observed,

smoothed and predicted LE for the population of men, women and both sexes during 2002-2011 are plotted in the Fig.1 to Fig.3 respectively.




Fig. 1: Observed, Smoothed and Predicted LE for Male Population of Urban Area in Bangladesh During 2002-2011. X: Years and Y: LE.




Fig. 2: Observed, Smoothed and Predicted LE for Female Population of Urban Area in Bangladesh During 2002-2011. X: Years and Y: LE.




Fig. 3: Observed, Smoothed and Predicted LE for Both Sexes of the Population of Urban Area in Bangladesh During 2002-2011. X: Years and Y: LE.

The model, that is, two parameters linear regression model is constructed to LE for the population of male, female and both sexes of urban area in Bangladesh and the constructed models are as follows:

$$y_t = -259.330 + 0.163t \text{ for male} \quad (1)$$

$$\text{t-stat} \quad (-9.7387) \quad (12.2845)$$

$$\text{P-value} \quad (0.0000) \quad (0.0000)$$

$$y_t = -620.676 + 0.344t \text{ for female} \quad (2)$$

$$\text{t-stat} \quad (-9.6447) \quad (10.7143)$$

$$\text{P-value} \quad (0.0000) \quad (0.0000)$$

$$y_t = -378.084 + 0.222t \text{ for both sexes} \quad (3)$$

$$\text{t-stat} \quad (-8.212) \quad (9.693)$$

$$\text{P-value} \quad (0.0000) \quad (0.0000)$$

The enumerated CVPP, ρ_{cv}^2 related to their R^2 are demonstrated in Table 3. From the table it is observed that the constructed models are highly cross-validated and their shrinkages coefficients are very lowest. In the table, it is found that the constructed models (1) – (3) are stable more than 92%, 90% and 91% respectively. Moreover, all the parameters of the constructed models (1) – (3) are significant explaining large proportion of variance. The stabilities of R^2 of these constructed models are more than 97%.

Table 3: Information on Model Fittings and Estimated CVPP of the Predicted Equations

| Models | N | k | R^2 | ρ_{cv}^2 | Shrinkage | Variance explained (%) |
|------------|----|---|---------|---------------|------------|------------------------|
| Equation 1 | 10 | 1 | 0.94966 | 0.928805 | 0.02085514 | 94.966 |
| Equation 2 | 10 | 1 | 0.93485 | 0.907859 | 0.02699071 | 93.485 |
| Equation 3 | 10 | 1 | 0.92154 | 0.889035 | 0.032505 | 92.15 |

From these statistics of Table 4, it is also decided that all these fitted models are highly significant. For this reason, the fits of all these models are good. Then, the projected values of LE are estimated by applying these constructed regression models that are exposed in Table 5. After plotting these

LE of men, women and both sexes of the population of urban area during 2012-2051 in the Fig.4, it shows an upward trend due to time like the trend during 2002-2011. These projections of LE give an idea regarding the future size of the oldest population in Bangladesh.

Table 4: Calculated and Tabulated Values of F-statistics of the Predicted Equations

| Models | n | K | Cal. F | Tab.F (at 1% level) |
|------------|----|---|--------|----------------------|
| Equation 1 | 10 | 2 | 150.92 | 11.3 with (1,8) d.f. |
| Equation 2 | 10 | 2 | 114.79 | 11.3 with (1,8) d.f. |
| Equation 2 | 10 | 2 | 93.96 | 11.3 with (1,8) d.f. |

Table 5: Projected LE for Male, Female and Both Sexes of the Population of Urban Area in Bangladesh during 2012-2051

| Year | Male | Female | Both Sexes |
|------|------|--------|------------|
| 2012 | 68.6 | 70.7 | 69.4 |
| 2013 | 68.8 | 71.1 | 69.6 |
| 2014 | 69.0 | 71.4 | 69.9 |
| 2015 | 69.1 | 71.8 | 70.1 |
| 2016 | 69.3 | 72.1 | 70.3 |
| 2017 | 69.4 | 72.4 | 70.5 |
| 2018 | 69.6 | 72.8 | 70.7 |
| 2019 | 69.8 | 73.1 | 71.0 |
| 2020 | 69.9 | 73.5 | 71.2 |
| 2021 | 70.1 | 73.8 | 71.4 |
| 2022 | 70.3 | 74.2 | 71.6 |
| 2023 | 70.4 | 74.5 | 71.8 |
| 2024 | 70.6 | 74.8 | 72.1 |
| 2025 | 70.7 | 75.2 | 72.3 |
| 2026 | 70.9 | 75.5 | 72.5 |
| 2027 | 71.1 | 75.9 | 72.7 |
| 2028 | 71.2 | 76.2 | 72.9 |
| 2029 | 71.4 | 76.6 | 73.2 |
| 2030 | 71.6 | 76.9 | 73.4 |
| 2031 | 71.7 | 77.2 | 73.6 |

| | | | |
|------|------|------|------|
| 2032 | 71.9 | 77.6 | 73.8 |
| 2033 | 72.0 | 77.9 | 74.0 |
| 2034 | 72.2 | 78.3 | 74.3 |
| 2035 | 72.4 | 78.6 | 74.5 |
| 2036 | 72.5 | 79.0 | 74.7 |
| 2037 | 72.7 | 79.3 | 74.9 |
| 2038 | 72.9 | 79.7 | 75.1 |
| 2039 | 73.0 | 80.0 | 75.4 |
| 2040 | 73.2 | 80.3 | 75.6 |
| 2041 | 73.4 | 80.7 | 75.8 |
| 2042 | 73.5 | 81.0 | 76.0 |
| 2043 | 73.7 | 81.4 | 76.2 |
| 2044 | 73.8 | 81.7 | 76.5 |
| 2045 | 74.0 | 82.1 | 76.7 |
| 2046 | 74.2 | 82.4 | 76.9 |
| 2047 | 74.3 | 82.7 | 77.1 |
| 2048 | 74.5 | 83.1 | 77.3 |
| 2049 | 74.7 | 83.4 | 77.6 |
| 2050 | 74.8 | 83.8 | 77.8 |
| 2051 | 75.0 | 84.1 | 78.0 |




Fig. 4: Projected LE for Male, Female and Both Sexes of the Population of Urban Area in Bangladesh during 2012-2051. X: Years (Time) and Y: Projected LE.

DISCUSSION

In the current study, the core objective was to project the LE of the urban populations of Bangladesh by sex. The simple linear regression model was employed as a method of projecting the LE. The projections were accomplished up to the year 2051. The results revealed that the LE of the urban populations of Bangladesh by sex increase in accordance with time and female tend to live longer than male and it clearly exhibited the differences in LE between male and female. These findings conform to the study of Malaysian population (Islam et al., 2017). In this study, actually the LE was treated as the average number of years of a person who can expect to live provided existing mortality patterns. The LE is treated as the most commonly used measurement of health status of population as well as, in fact, mortality. Nonetheless, LE should be familiar as a measure of length of life rather than quality of life, on the contrary, it does not count the full burden of illness and disability.

In 2002, the LE in the urban populations of Bangladesh was 67.2(male, 67; female, 67.3) years (BBS, 2013), on the other hand, by 2051, it would be increased as 78 (male, 75; female, 84.1) years (projected). The long life can be attributed by a number of various types of factors together with rising living standards, enhanced lifestyle and improved quality education, as well as greater access to quality as well as available health services which are significantly increased LE in

Bangladesh, especially in the urban area of Bangladesh. Therefore, the demographic and socioeconomic changes, and availability of health facilities persuade LE (Mondal and Shitan, 2013, 2014; Mondal et al, 2015). In Bangladesh, the expansion of population and the demographic transition since the 1971 were accompanied by major socioeconomic development. LE is affected by different factors such as socioeconomic status, social factors, genetic factors and environmental factors like income, education, the quality of the health system, health behaviors such as tobacco and alcohol consumption, poor nutrition, overcrowded housing, lack of clean drinking water, adequate sanitation, and lack of exercise.

After the long years of socioeconomic development, Bangladesh had transformed into middle income country. Accordingly, the socioeconomic development determines the improvements in social sectors and increases LE. Increases in LE have also been attributed by the substantial improvements in sanitation and the availability as well as easy access to clean drinking water (Mondal and Shitan, 2015). The sanitation system in Bangladesh has been improving significantly which is imperatively as well as significantly high as the developing nations. An increase in LE was driven mainly by the enhancement in sanitation in the time of the nineteenth and early twentieth century's (Oeppen and Vaupel, 2002). Environmental quality is a very important factor affecting health and morbidity: air and water pollution, depletion of natural resources, soils deterioration is all capable of increasing human mortality that mean reducing life expectancy (Elo and Preston, 1992; Pope, 2000; Pope et al., 2004 and Evans and Smith, 2005)

In deaths and LE, the sex differences vary from country to country or any region to region. But, in most countries, generally male live shorter lives than that of female, sometimes by a margin of as much as 5-10 years. This study was investigated that the LE for female would be higher compared to male in Bangladesh. The deaths for women were found lower than that of men for all ages. In general, more boys than girls die in infancy period and during each subsequent years of life, mortality for male exceed than that of female (Islam, 2003). As a consequence, the gender difference would be more broadened in the future of the country.

Conclusions

In this paper, it is observed that LE for the population of urban area in Bangladesh follow a simple linear regression model. It is investigated that the significance of these constructed models is very

soaring because of test statistics-F. Then, LE for the population of male, female and both sexes of urban area in Bangladesh are projected using these models during 2012-2051. It is also observed that people in Bangladesh tend to have substantial longer life expectancy rate. These might be used as predicted LE for the population of male, female and both sexes of urban area in Bangladesh for 2012-2051 for further higher study. The LE for the population of male, female and both sexes of urban area in Bangladesh in 2051 will be 75, 84.1 and 78 years respectively which exhibits that the urban population of Bangladesh are radically tending to be old aged population. Proper planning for the allocation of public and private resources may be taken based on the findings of this study for the better lifestyle of urban residents. More research study should be needed in this arena.

REFERENCES

- BBS. 2013. Statistical Year Book of Bangladesh 2012. Government of the People's Republic of Bangladesh: Bangladesh Bureau of Statistics (BBS).
- BBS. 2014. Statistical Year Book of Bangladesh 2013. Government of the People's Republic of Bangladesh: Bangladesh Bureau of Statistics (BBS).
- Chiang L. 1968. The Life Table and Its Construction, Introduction to Stochastic Processes in Biostatistics, New York. *John Wiley and Sons* 189–214.
- Elo IT, Preston SH. 1992. Effects of Early-Life Conditions on Adult Mortality: A Review. *Population Index*. 58 (2): 186-212.
- Evans MF, Smith VK. 2005. Do New Health Conditions Support Mortality-Air Pollution Effects? *Journal of Environmental Economics and Management*. 50: 496-518.
- Gujarati N. 1998. Basic Econometrics, Third Edition, *McGraw Hill, Inc.*, New York.
- Hossain MS, Islam MR. 2013. Age Specific Participation Rates of Curacao in 2011: Modeling Approach. *American Open Computational and Applied Mathematics Journal*. 1(2): 08 - 21.
- Islam MR. 2003. Modeling of Demographic Parameters of Bangladesh-An Empirical Forecasting, *Unpublished Ph.D. Thesis*, Rajshahi University.
- Islam MR. 2004. Mathematical Modeling of Age Specific Fertility Rates in Urban Area of Bangladesh. *J. of Science, Rajshahi University Studies, Part- B*. 32: 229-240.
- Islam MR, Islam N, Rahman M. 2004. Trends and Patterns of Reproductivity Decline in Bangladesh: 1956-1998. *Man in India*. 84(1 & 2): 85-97.

- Islam MR. 2005a. Construction of Abridged Life Tables and Indirect Estimation of Some Mortality Measures of Bangladesh in 2005. *Journal of Population*. 11(2): 117-130.
- Islam MR. 2005b. Mathematical Modeling of Age Specific Marital Fertility Rates in Urban Area of Bangladesh. *Pakistan J. of Statistics*. 21(3): 289-295.
- Islam MR. 2006a, Modeling of Age Structure and ASDRs for Both Sexes of Bangladesh. *J. of Social Anthropology*. 3(1): 51-56.
- Islam MR. 2006b. Modeling ASFRs of Bangladesh: An Application of Polynomial Model. *South Asian Anthropologist*. 6(2): 97-100.
- Islam MR. 2006c. Age Associated with Expectation of Life (ex) for Male of Bangladesh: Application of Polynomial Model. *J. of Engineering and Applied Sciences*. 1(2): 161-164.
- Islam MR. 2007a. Construction of Abridged Life Tables and Indirect Estimation of Some Mortality Measures of Bangladesh in 2005. *Journal of Population*. 11(2): 117-130.
- Islam MR. 2007b. Predicting Population for Male of Rural Area in Bangladesh. *Journal of Statistical Research of Iran*. 4(2): 227-238.
- Islam MR. 2007c. Estimation of Some Mortality Measures, Modeling of l_x and Age Associated with Force of Mortality of Bangladesh in 2006. *Middle East J. of Age and Ageing*. 4(5): 23-28.
- Islam MR. 2008. Modeling Age Structure for Female Population in Rural Area of Bangladesh. *Journal of Interdisciplinary Mathematics*. 11(1): 29-37.
- Islam MR. 2011. Modeling of Diabetic Patients Associated with Age: Polynomial Model Approach. *International Journal of Statistics and Applications*. 1(1): 1-5.
- Islam MR. 2012a. Mathematical Modeling of Age and of Income Distribution Associated with Female Marriage Migration in Rajshahi, Bangladesh. *Research Journal of Applied Sciences, Engineering and Technology*. 4(17): 3125-3129.
- Islam MR. 2012b. Modeling and Projecting Population for Muslim of Urban Area in Bangladesh. *International Journal of Probability and Statistics*. 1(1): 4-10.
- Islam MR. 2013. Modeling Age Structure and ASDRs for Human Population of Both Sexes in Bangladesh. *International Journal of Anthropology*. 28(1): 47-53.
- Islam MR, Hossain MS. 2013a. Mathematical Modeling of Age Specific Adult Literacy Rates of Rural Area in Bangladesh. *American Open Demography Journal*. 1(1): 01-12.

Islam MR, Hossain MS. 2013b. Mathematical Modeling of Age Specific Participation Rates in Bangladesh, *International Journal of Scientific and Innovative Mathematical Research*. 1(2): 150-159.

Islam MR, Ali K, Islam N. 2013. Construction of Life Table and Some Mathematical Models for Male Population of Bangladesh. *American Journal of Computational and Applied Mathematics*. 3(6): 269-276.

Islam MR, Hossain MS. 2014a. Some Models Associated with Age Specific Adult Literacy Rates of Urban Area in Bangladesh. *International Journal of Ecosystem*. 4(2): 66-74.

Islam MR, Hossain MS. 2014b. Mathematical Modeling of Age Specific Adult Literacy Rates in Bangladesh. *Advances in Life Sciences*. 4(3): 106-113.

Islam MR, Hossain MS, Faroque O. 2014. U-Shaped Pattern of Employees' Job Satisfaction: Polynomial Model Approach. *International Journal of Ecosystem*. 4(4): 170-175.

Islam MR, Hoque MN. 2015. Mathematical Modeling and Projecting Population of Bangladesh by Age and Sex from 2002 to 2031. Emerging Techniques in Applied Demography. *Applied Demography Series 4*. 5: 53-60.

Islam MR, Shitan M, Mondal MNI. 2017. Projection of Life Expectancy at Birth of Malaysian Population: A Modelling Approach. *Journal of The Indian Academy of Geriatrics*. 13: 68-72.

Lima M, Barros A, Cesar G, Goldbaum M, Carandina L, Ciconelli M. 2009. Health Related Quality of Life Among the Elderly: A Population-Based Study Using SF-36 Survey. *Cad Saude Publica*. 25(10): 2159-2167.

Islam MR, Khan MN, Hoque MN, and Khan MMA. 2020. Projection of Female Urban Population in Bangladesh. Chapter 22, Population Change and Public Policy, *Applied Demography Series 11*, 433-444, https://doi.org/10.1007/978-3-030-57069-9_22.

Islam M.R. and Shitan M. 2022. Modeling and Projection of Some Fertility Parameters of Malaysian Population, *Human Biology Review*. 11(2): 66-78.

Mondal MNI, Shitan M. 2014. Relative Importance of Demographic, Socioeconomic and Health Factors on Life Expectancy in Low- and Lower-Middle-Income Countries. *J Epidemiol*. 24(2): 117-124.

- Mondal MNI, Shitan M. 2013. Impact of Socio-Health Factors on Life Expectancy in the Low and Lower Middle Income Countries. *Iranian J Publ Health*. 42(12):1354-1362.
- Mondal MNI, Ullah MMMN, Islam MR, Rahman MS. 2015. Socio-demographic and Health Determinants of Inequalities in Life Expectancy in Least Developed Countries. *International Journal of MCH and AIDS*. 3(2): 96-105.
- Mondal MNI, Shitan M. 2015. Author's Response: Regarding the Relative Importance of Demographic, Socioeconomic and Health Factors on Life Expectancy in Low- and Lower-Middle-Income Countries. *J Epidemiol*. 25(6): 460.
- Oeppen J, Vaupel JW. 2002. Broken Limits to Life Expectancy. *Science*. 296: 1029.
- Pope CA, III. 2000. Epidemiology of Fine Particulate Air Pollution and Human Health: Biologic Mechanisms and Who's at Risk? *Environmental Health Perspectives*. 108(4): 713-723.
- Pope CA, III, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, Godleski JJ. 2004. Cardiovascular Mortality and Long term Exposure to Particulate Air Pollution: Epidemiological Evidence of General Pathophysiological Pathways of Disease. *Circulation*. 109(1): 71-77.
- Parahyba I, Veras R, Melzer D. 2005. Disability among Elderly Women in Brazil. *Rev Saude Publica*. 39(3): 383-391.
- Stevens J. 1996. Applied Multivariate Statistics for the Social Sciences, Third Edition, Lawrence Erlbaum Associates, Inc., Publishers, New Jersey.
- Szwarchwald L, Mota C, Damacena N, Pereira S. 2011. Health Inequalities in Rio de Janeiro, Brazil: Lower Healthy Life Expectancy in Socioeconomically Disadvantaged Areas. *Am J Public Health*. 101(3): 517-523.
- Velleman F. 1980. Definition and Comparison of Robust Nonlinear Data Smoothing Algorithms. *Journal of the American Statistical Association*. 75(371): 609-615.
- WHO. 2013. World Health Statistics 2013, World Health Organization. Website accessed March 2014.
- Zunzunegui V, Alvarado E, Beland F, Vissandjee B. 2009. Explaining Health Differences Between Men and Women in Later Life: A Cross-City Comparison in Latin America and the Caribbean. *Soc Sci Med*. 68: 235-242.